

IV) 病虫害発生程度別面積

農作物名	作付面積	病虫害名	程度別発生面積					防除面積		被害面積	被害量	平年発生面積	備考
			甚	多	中	少	計	実	延				
イネ	20,600 ^{ha}	苗立枯病	1	9	75	85	19,300	22,200	10	t	111.7		
		苗葉いもち	2	48	50	0	0	0	2		70.6		
		穂いもち	5	100	535	640	930	11,400	105		13,402.0		
		紋いもち	5	25	500	530	12,010	13,040	300		10,302.0		
		白馬鹿病	10	160	1,730	12,200	14,100	11,600	12,400	1,900	15,063.0		
		イネシンガレセンチュウ	5	85	460	550	18	18	90		832.5		
		マシロ病		2	55	57	19,200	19,200	2		236.0		
		イネシロ病	60	180	460	8,300	9,000	(同)19,000	(同)19,000	1		9.1	
		イネシロ病		2	13	15	(同)6,100	(同)6,100	700		7,250.0		
		イネシロ病		2	13	15	(同)2,500	(同)2,500	2		362.5		
		イネシロ病		3	3	6	(同)13,000	(同)15,000	0		7,470.0		
		イネシロ病		3	3	6	(同)13,000	(同)15,000	3		10.9		
		イネシロ病		400	2,000	17,000	19,400	(同)13,000	(同)15,000	400		20,475.5	
		イネシロ病		6	6	6	0	0	0		9.8		
		イネシロ病		0	0	0	0	0	0		0		
		イネシロ病		3	127	1,900	2,030	(同)400	(同)400	130		3,053.0	
		イネシロ病		1	9	10	180	180	1		11.1		
		イネシロ病		5	20	600	625	7	7	25	433.0		
		イネシロ病		20	1,000	1,020	1,100	1,100	20		1,404.5		
		イネシロ病		10	1,500	1,510	(同)13,000	(同)15,000	10		1,816.0		
		イネシロ病		1,000	15,010	16,010	(同)13,000	(同)15,000	400		19,926.5		
		イネシロ病		20	50	71	(同)13,000	(同)15,000	1		6,363.0		
		イネシロ病		1	100	1,200	1,301	(同)13,000	(同)13,000	10		1,340.0	
		イネシロ病		40	40	40	0	0	0		40.0		
		イネシロ病		100	500	600	(同)13,000	(同)13,000	90		3,529.5		
		イネシロ病		2,000	2,000	2,000	(同)13,000	(同)13,000	10		6,066.0		
		イネシロ病		20	100	122	125	125	2		89.6		
		イネシロ病	5	2	3,000	12,500	15,507	13,000	15,000	1,200	11,497.8		
		イネシロ病		100	100	600	750	50	50	25	1,605.9		
		イネシロ病		50	100	600	750	50	50	20	5,157.0		
		イネシロ病		50			50	50	0	0	275.0		
		イネシロ病					0	0	0	0	0.7		
		イネシロ病	5		500	2,000	2,505	1,500	1,500	50	8,330.0		
		イネシロ病		1,000	2,000	8,500	11,500	13,000	13,000	50	18,912.0		

農作物名	作付面積	病 害 虫 名	程 度 別 発 生 面 積					防 除 面 積		被害面積	被害量	平年発 生面積	備 考				
			甚	多	中	少	計	実	延								
ム ギ	498 ^{ha}	赤小 黄黒 う赤 雪黒 黒縞 斑 さ さ さ さ ど か 腐 穂 節 萎 葉 び び び び び こ び 縮	病	ha	ha	ha	ha	ha	ha	ha	ha	t	ha				
			病		1	3	33	37	0	0	4	13.8					
			病				20	20	0	0	0	30.9					
			病					0	0	0	0	0					
			病					0	—	—	0	0					
			病				13	13	(同)	370	(同)	370	23.7				
			病				13	13		450		900	28.1				
			病					0		0		0	0				
			病				1	1		0		0	5.7				
			病				7	7		—		—	13.4				
			病				1	1		—		—	9.3				
			病				7	8		0		0	23.9				
			ダ イ ズ	1,240	紫さ 菌立 炭枯 べ 葉 モ 萎 ア ハ カ 斑 核 性 疽 と 焼 イ 縮 ク シ ト ヨ シ 病 害 病 病 病 病 病 類 ウ 類	病		3	12	185	200	170	180	15	325.0		
病								1	1	0	0	0	0.9				
病								1	1	0	0	0	1.5				
害							1	4	5	0	0	1	8.8				
病								4	4	0	0	0	5.8				
病							5	115	120	—	—	40	352.0				
病							5	15	300	320	0	0	20	281.0			
病							5	5	70	75	—	—	5	98.0			
病									0	0	—	—	0	0			
類							5	50	200	255	(同)	400	(同)	500	30	353.5	
類							4	70	200	450	724	400	600	170	722.0		
類							4	40	200	550	794	(同)	400	(同)	500	160	716.1

農作物名	作付面積	病 害 虫 名	程度別発生面積					防除面積		被害面積	被害量	平年発 生面積	備 考	
			甚	多	中	少	計	実	延					
ナ シ	116 ha	黒斑病	ha	ha	ha	ha	ha	ha	ha	ha	ha	t	ha	
		黒星病		6	19	38	63	(同)	63	(同)	819	25		102.9
		赤星病			1	13	14	(同)	116	(同)	696	1		42.8
		ナシヒメシクイガ	1	4	15	12	12	(同)	116	(同)	348	0		45.8
		モシクイガ		2	8	10	20	(同)	115	(同)	230	6		12.7
		ナシオシクイガ				2	2	(同)	115	(同)	115	4		19.3
		ハマキムシ				20	20	(同)	115	(同)	230	0		1.3
		ハマダニ		20	30	25	75	(同)	115	(同)	345	47		29.5
		クワコナカイガラムシ				15	15	(同)	115	(同)	20	2		112.3
		カメムシ		10	20	25	55	(同)	115	(同)	115	50		17.4
		ナシチビガ				10	10	(同)	115	(同)	115	0		39.4
		アブラムシ	1	6	40	60	107	(同)	115	(同)	230	44		1.5
		吸蛾類				10	10	(同)	115	(同)	230	2		118.5
		クワゴマダラヒトリ				1	1	(同)	115	(同)	5	0		14.5
		ナシグンバライ				10	10	(同)	115	(同)	115	0		1.5
		ニセナシサビダ		10	20	25	55	(同)	115	(同)	345	5		9.0
		ナシホソ				5	5	(同)	115	(同)	115	0		44.6
												9.5		
ブ ド ウ	341	晩腐病			6	22	28	(同)	175	(同)	349	6		19.4
		さび病				9	9	(同)	279	(同)	558	0		98.8
		褐斑病		4	26	40	70	(同)	279	(同)	558	30		146.6
		黒とう病			4	16	20	(同)	87	(同)	174	4		11.3
		べと病		7	15	52	74	(同)	279	(同)	558	22		162.3
		灰色かび病			13	64	77	(同)	279	(同)	418	13		55.0
		うど病				7	7	(同)	63	(同)	126	0		11.2
		ブドウスカシバリ				30	30	(同)	340	(同)	340	0		53.0
		ブドウトラカミキリ				100	100	(同)	340	(同)	340	33		191.5
		フタテンヒメヨコバイ			20	160	180	(同)	340	(同)	340	3		219.0
		チャノキイロアザミウマ	1	1	30	100	132	(同)	340	(同)	340	30		122.1
		コガネムシ			4	26	30	(同)	10	(同)	10	2		19.6
		ハマキムシ				50	50	(同)	55	(同)	100	19		76.0
		ハマダニ	1	4	45	110	160	(同)	340	(同)	340	35		144.0
		クワゴマダラヒトリ					0	(同)	0	(同)	0	0		0.0
		カイガラムシ			20	45	65	(同)	100	(同)	100	27		73.0

農作物名	作付面積	病 害 虫 名	程 度 別 発 生 面 積					防 除 面 積		被害面積	被害量	平年発 生面積	備 考	
			甚	多	中	少	計	実	延					
カ キ	563 ^{ha}	炭疽病	ha	ha	ha	ha	ha	ha	ha	ha	t	ha		
		うどんこ病			3	25	28	(同)	450	(同) 1,800	3		53.6	
		うどんこ病			18	50	68	(同)	450	(同) 1,800	18		258.0	
		円形星状病	20	47	55	57	179		450	1,350	122		171.8	
		角斑病		10	13	20	43		450	1,350	23		32.5	
		黒星病			10	17	27		370	370	10		18.3	
		カキミガシ				70	70	(同)	300	(同) 900	14		72.5	
		フジコナカイガラムシ	3	10	30	55	98		500	1500	43		46.6	
		カメムシ類			20	30	50		500	500	51		221.5	
		チャノキイロアザミウマ	2	5	20	20	47	(同)	500	(同) 1,000	15		144.5	
		カキクダアザミウマ					110	(同)	500	(同) 1,000	0		164.1	
		吸蛾類					0		0	0	0		0.0	
		クワゴマダラヒトリ					100		500	500	0		0.1	
ハマキムシ類			10	110	120	(同)	500	(同) 1,000	34		159.5			

農作物名	作付面積	病虫害名	程度別発生面積					防除面積		被害面積	被害量	平年発生面積	備考	
			甚	多	中	少	計	実	延					
春夏 メロン	72 ha	べつと	ha	ha	ha	ha	ha	ha	ha	ha	ha	ha	ha	
		るる			<1	2	2	51	160	<1	t	8.6		
		枯割			2	10	12	51	160	2		6.9		
		斑根			<1	<1	<1	<1	<1	<1		<1		
		点腐			<1	2	5	13	13	1		4		
		病病			<1	3	3	6	6	<1		3		
秋冬 ダイコン	302 ha	モア												
		ブ		3	7	40	50	70	210	2		57.1		
		ラ		3	15	42	61	70	210	7		63.4		
		ム			1	7	8	50	60	1		17.0		
		シ				1	85	86	—	—	1		114.7	
		病類			15	112	130	150	230	3		155.0		
秋冬 ハクサイ	182 ha	モ												
		軟												
		白												
		黒		1	11	75	91	100	300	12		124.7		
		ア						(同)	100	(同)	1		35.4	
		病類												
秋冬 ハクサイ	182 ha	ブ												
		ラ		1	7	73	81	150	230	1		128.0		
		ム						(同)	170	(同)	2		26.2	
		シ		1	5	24	30	70	100	1		33.0		
		病類												
		シ		1	5	24	30	70	100	1		33.0		
秋冬 ハクサイ	182 ha	ダ												
		イ												
		コ												
		ナ												
		ハ												
		病類												
秋冬 ハクサイ	182 ha	タ												
		マ												
		ナ												
		シ												
		ロ												
		病類												
秋冬 ハクサイ	182 ha	モ												
		ス		2	2	18	20	(同)	170	(同)	0		40.0	
		モ		8	25	115	150	(同)	170	(同)	12		120.0	
		ン												
		シ												
		病類												
秋冬 ハクサイ	182 ha	ハ												
		イ		2	3	15	20	30	60	2		9.4		
		マ												
		ダ												
		ラ												
		病類												

農作物名	作付面積	病 害 虫 名	程 度 別 発 生 面 積					防 除 面 積		被害面積	被害量	平年発 生面積	備 考	
			甚	多	中	少	計	実	延					
夏秋 キャベツ	ha 62	黒菌 腐病	ha	ha	ha	ha	ha	ha	ha	ha	ha	t	ha	
		株核病			1	12	13	50	125	1			37.8	
		モシロウ			2	12	14	50	75	2			20.6	
		コシロウ			2	18	20	—	—	2			19	
		ヨトウムシ		1	5	8	13	(同) 60	(同) 240	1			19.8	
		ハスモンヨトウ		1	5	20	26	(同) 60	(同) 240	5			53.0	
		アブラムシ	2	3	2	7	10	(同) 60	(同) 100	1			11.1	
		タマナギン		3	15	20	40	(同) 60	(同) 240	8			38.8	
		ウワバ		1	3	12	15	55	80	1			27.3	
冬キャベツ	108	黒菌 腐病				2	20	22	90	180	2		34.2	
		株核病				5	15	20	90	180	5		22.7	
		モシロウ				7	23	30	—	—	7		37	
		コシロウ				1	19	20	(同) 100	(同) 200	0		40.5	
		ヨトウムシ		1	5	25	30	(同) 100	(同) 200	1			54.0	
		ハスモンヨトウ	1	6	3	15	19	(同) 100	(同) 200	1			19.6	
		アブラムシ		1	18	65	90	(同) 100	(同) 250	7			56.4	
		タマナギン		1	1	18	20	90	110	1			32.5	
		シロイチモ			2	15	17	(同) 100	(同) 150	0			32.4	
タマネギ	151	白色疫病				10	10	130	380	0		16.0		
		ボトリチス	1	5	10	47	63	130	460	16		74.7		
		葉枯れ				40	40	—	—	—		92.8		
イチゴ	27	灰色かんび		1	2	18	21	27	108	21		23.3		
		うど		1	3	14	18	27	270	18		22.0		
		炭疽				4	4	—	—	—		—		
		アブラムシ		1	3	17	21	24	60	2		28.3		
		ハダニ		1	5	20	26	25	75	3		33.1		
キク	29	ハスモンヨトウ		3	10	12	25	—	—	3		16.5		
		アブラムシ			1	3	4	29	150	0		4.3		
		アザミ			1	5	6	29	160	1		6.0		